

Table I: Age-Adjusted and Unadjusted Prevalence* of Selected Chronic Conditions among Adults, by County, Oregon 2004-2007

County	Arthritis		Asthma		Heart Attack		Coronary Heart Disease	
	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted
OREGON		26.9%		9.9%		3.5%		3.6%
Baker	39.5%	34.4%	9.0%	9.1%	3.8%	2.6%	6.0%	4.3%
Benton	23.9%	27.4%	8.4%	9.1%	2.7%	3.2%	3.0%	3.3%
Clackamas	26.5%	25.8%	10.4%	10.4%	3.1%	2.9%	3.7%	3.5%
Clatsop	30.2%	28.5%	9.8%	10.0%	5.7%	4.9%	6.2%	5.5%
Columbia	31.5%	30.5%	11.1%	11.0%	5.6%	5.2%	3.5%	3.1%
Coos	35.1%	28.7%	9.5%	9.3%	4.7%	3.5%	6.0%	4.5%
Crook	29.8%	27.2%	9.9%	10.2%	4.4%	3.8%	5.4%	4.6%
Curry	36.3%	27.2%	11.5%	9.9%	9.1%	5.4%	11.2%	8.4%
Deschutes	30.1%	28.4%	8.8%	9.0%	3.6%	3.2%	3.5%	3.2%
Douglas	39.2%	35.7% **	12.6%	12.8% **	6.5%	5.6% **	6.7%	5.4% **
Grant	25.0%	20.7%	14.8%	17.2%	6.0%	5.7%	3.1%	2.2%
Harney	35.3%	31.2%	8.5%	8.0%	5.1%	4.2%	4.7%	4.0%
Hood River	24.2%	22.4%	5.7%	5.9%	4.6%	4.5%	4.3%	4.1%
Jackson	29.0%	26.4%	9.3%	9.2%	3.3%	2.6%	3.7%	3.1%
Jefferson	31.4%	30.0%	7.2%	7.0%	4.2%	3.7%	3.4%	2.8%
Josephine	33.6%	29.0%	11.3%	12.4%	4.8%	3.5%	5.5%	4.3%
Klamath	39.2%	36.4%	8.7%	8.8%	6.7%	5.7% **	5.7%	4.9%
Lake	45.3%	37.0%	6.4%	6.4%	1.4%	1.0%	1.5%	1.0% **
Lane	28.8%	28.4%	10.0%	10.0%	3.2%	3.1%	3.4%	3.2%
Lincoln	34.7%	29.6%	11.9%	12.1%	8.0%	6.4% **	6.0%	5.3%
Linn	33.2%	31.0%	11.2%	11.5%	4.9%	4.3%	3.7%	3.2%
Malheur	25.3%	24.4%	6.8%	6.9%	1.7%	1.7%	3.0%	2.9%
Marion	27.3%	27.3%	10.0%	10.0%	4.0%	4.0%	3.6%	3.6%
Morrow	20.3%	20.0%	9.6%	10.0%	2.4%	2.3%	8.1%	8.2%
Multnomah	24.5%	25.6%	9.9%	9.9%	3.2%	3.4%	3.3%	3.6%
Polk	23.0%	22.0%	8.6%	8.7%	4.3%	3.9%	4.9%	4.4%
Tillamook	33.2%	27.2%	8.4%	7.8%	4.6%	3.8%	4.9%	3.8%
Umatilla	32.6%	31.9%	9.0%	9.0%	4.0%	3.8%	4.2%	4.0%
Union	29.7%	29.3%	10.9%	10.9%	3.3%	3.0%	5.9%	5.1%
Wallowa	25.1%	19.0%	8.9%	8.8%	3.5%	2.3%	5.9%	3.8%
Washington	21.4%	23.1% **	9.1%	9.2%	2.1%	2.3% **	2.5%	2.8%
Yamhill	26.3%	27.2%	11.4%	11.2%	4.4%	4.5%	3.7%	3.8%
Gilliam/Wheeler	27.2% †	16.4% †	4.8%	3.9%	4.6% †	2.5% †	4.0% †	2.2% †
Sherman/Wasco	27.1%	22.6%	7.7%	7.3%	3.8%	3.0%	4.2%	3.1%

*Unadjusted rates provide a description of the absolute burden of disease or risk factor for a county. Age-adjusted rates allow you to compare rates for a given county to the State's rate. When comparing counties with Oregon or with each other, it's best to use age-adjusted rates.

** Statistically significant difference compared to Oregon.

† % based on less than 50 respondents or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county. Age-adjusted to the 2000 Standard Population using 3 age groups (18-34, 35-54, and 55+).

Sources: BRFSS 2004-2007

Table I (continued): Age-Adjusted and Unadjusted Prevalence* of Selected Chronic Conditions among Adults, by County, Oregon 2004-2007

County	Stroke		Diabetes		High Blood Pressure		High Blood Cholesterol	
	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted	unadjusted	age-adjusted
OREGON		2.3%		6.5%		24.8%		32.3%
Baker	1.3%	0.9%	9.8%	8.4%	42.5%	38.8%**	51.3%	52.0%**
Benton	1.7%	2.0%	3.7%	4.3%**	21.3%	24.1%	33.9%	32.6%
Clackamas	2.4%	2.3%	5.6%	5.4%**	24.6%	23.9%	36.6%	32.3%
Clatsop	3.6%	2.9%	9.0%	8.1%	27.1%	24.4%	41.8%	34.3%
Columbia	1.9%	1.7%	8.9%	8.4%	31.3%	28.8%	31.5%	27.4%
Coos	2.8%	2.2%	8.8%	6.8%	37.6%	32.7%**	35.7%	29.8%
Crook	1.9%	1.5%	5.0%	4.3%	22.8%	21.0%	31.9%	31.3%
Curry	4.6%	3.0%	9.4%	6.9%	42.5%	35.0%	37.5%	26.6%
Deschutes	2.2%	1.9%	5.5%	5.1%	25.3%	23.8%	39.4%	33.8%
Douglas	4.3%	3.7%**	9.8%	8.5%**	34.2%	30.2%	41.9%	36.4%
Grant	6.1%	5.8%	5.4%	3.8%	29.9%	22.9%	35.7%†	29.4%†
Harney	4.9%	3.2%	9.3%	8.2%	31.1%	26.7%	44.5%†	43.1%†
Hood River	0.6%	0.6%	3.2%	3.0%**	19.3%	18.6%	34.3%	25.7%
Jackson	3.1%	2.5%	7.3%	6.7%	28.4%	26.0%	40.8%	35.2%
Jefferson	2.5%	2.1%	8.5%	7.9%	22.5%	20.0%	27.2%	22.8%
Josephine	2.7%	2.2%	7.7%	6.0%	30.4%	26.1%	44.6%	39.9%
Klamath	2.9%	2.5%	8.6%	7.9%	29.9%	27.2%	36.2%	30.9%
Lake	0%	0%	5.8%	4.8%	39.3%	32.9%	33.5%	23.7%
Lane	2.4%	2.4%	6.1%	6.0%	26.3%	25.8%	36.0%	31.6%
Lincoln	3.0%	2.2%	10.9%	8.7%	34.9%	30.5%	42.3%	32.2%
Linn	4.0%	3.7%**	8.8%	7.9%	31.1%	29.2%	34.6%	31.2%
Malheur	3.5%	3.2%	8.6%	8.3%	22.7%	22.4%	45.4%	43.8%
Marion	2.5%	2.5%	7.1%	7.2%	22.3%	22.5%	30.9%	27.3%**
Morrow	3.3%	3.4%	8.5%	8.5%	28.3%	28.1%	42.7%†	50.9%†
Multnomah	1.8%	1.9%	6.4%	6.7%	20.9%	22.0%	34.7%	32.7%
Polk	2.3%	2.1%	6.3%	5.8%	25.1%	23.8%	33.1%	28.8%
Tillamook	2.4%	1.6%	8.1%	6.3%	35.4%	28.4%	33.9%	21.8%**
Umatilla	2.6%	2.5%	8.9%	8.7%	32.5%	31.9%**	44.6%	40.2%
Union	2.9%	2.5%	5.3%	5.2%	19.2%	17.4%**	37.4%	29.1%
Wallowa	3.5%	2.3%	3.3%	2.2%**	28.4%	27.9%	31.4%	20.9%**
Washington	1.7%	1.9%	5.4%	5.9%	22.7%	24.4%	32.6%	30.5%
Yamhill	1.9%	2.0%	7.6%	7.8%	27.1%	27.3%	39.7%	37.8%
Gilliam/Wheeler	4.3%†	2.4%†	8.9%	6.2%	25.8%†	14.8%†	38.8%†	27.8%†
Sherman/Wasco	2.0%	1.6%	8.7%	7.2%	27.8%	24.6%	43.3%	31.9%

*Unadjusted rates provide a description of the absolute burden of disease or risk factor for a county. Age-adjusted rates allow you to compare rates for a given county to the State's rate. When comparing counties with Oregon or with each other, it's best to use age-adjusted rates.

** Statistically significant difference compared to Oregon.

† % based on less than 50 respondents or fewer than 12 persons in one age group; may not accurately reflect behavior of entire county. Age-adjusted to the 2000 Standard Population using 3 age groups (18-34, 35-54, and 55+).

Sources: BRFSS 2004-2007