



Oregon  
Sexually Transmitted Disease Program  
Syphilis Case Data  
2004


# Early Syphilis Cases and Incidence Oregon, 1985-2004



# Early Syphilis Cases by Stage of Diagnosis Oregon, 2000-2004



# Early Syphilis Cases by County of Residence Oregon, 2004




# Early Syphilis in the most affected Counties Oregon, 2004


<b>County</b>	<b>Cases</b>	<b>% of State</b>	<b>Incidence*</b>
Multnomah	37	64%	5.4
Washington	6	10%	1.2
Malheur	4	7%	12.6
Total of 3 counties	47	81%	---

\*per 100,000 population


# Early Syphilis Cases by Gender Oregon, 2000-2004




# Early Syphilis Cases by Age Group Oregon, 2000-2004




# Early Syphilis Cases by Gender and Age Oregon, 2004




# Early Syphilis Cases by Race Oregon 2000-2004




# Early Syphilis Cases by Race & Ethnicity Oregon, 2004




# Early Syphilis Cases among Men by MSM Status Oregon 2000-2004



# Early Syphilis Cases and Injection Drug Use Oregon 2000-2004



# Early Syphilis Cases Co-Morbidity with HIV and Gonorrhea Oregon, 2000-2004



Note: HIV co-morbidity not known for pre-2002 cases